




Fig. 5. Dissolution profiles of nifedipine solid dispersions (nifedipine: HPMC = 1:1) from matrix tablets with various ratios of lactose and dicalcium phosphate matrix granules (○: 1:0; ▽: 3:1; □: 1:1; △: 1:3; ◇: 0:1; ⊙: Adalat Retard®).

REFERENCES

1. Miller RA, Leung EM, Oates RJ. The compression of spheres coated with an aqueous ethylcellulose dispersion. *Drug Develop. Ind. Pharmac.* 1999;25:503-11.
2. Barra J, Falson-Rieg F, Doelker E. Modified drug release from inert matrix tablets prepared from formulations of identical composition but different organizations. *J. Control. Release* 2000;65:419-28.
3. Mallick S, Gupta BK, Ghosal SK. Assessment of bioavailability of experimental controlled release microcapsules of nifedipine. *Acta Poloniae Pharmac.* 2000;57:175-80.
4. Munoz-Ruiz A, Perales MCM, Antequera MV, Villar TP, Munoz N, Jimenez-Castellanos MR. Rheology and compression characteristics of lactose based direct compression excipients. *Int. J. Pharm.* 1995;95:201-7.
5. Patel NK, Upadhyay AH, Bergum JS, Reier GE. An evaluation of microcrystalline cellulose and lactose excipients using an instrumented single station tablet press. *Int. J. Pharm.* 1994;110:203-10.
6. Alderman DA. A review of cellulose esters in hydrophilic matrices for oral controlled-release dosage forms. *Int. J. Pharm. Tech. Prod. Manuf.* 1984;1-9.
7. Ford JL, Rubinstein MH, Hogan JE. Formulation of sustained release promethazine hydrochloride tablets using hydroxypropyl methylcellulose matrices. *Int. J. Pharm.* 1985;24:327-38.
8. Lin SY, Lin KH, Li MJ. Micronized ethylcellulose used for designing a directly compressed time-controlled disintegration tablet. *J. Control. Release* 2001;70:321-8.
9. Majid Khan G, Zhu JB. Ibuprofen release kinetics from controlled-release tablets granulated with aqueous polymeric dispersion of ethylcellulose II: Influence of several parameters and coexcipients. *J. Control. Release* 1998;56:127-34.
10. Wesseling M, Bodmeier R. Drug release from beads coated with an aqueous colloidal ethylcellulose dispersion, aquacoat, or an organic ethylcellulose solution. *Eur. J. Pharm. Biopharm.* 1999;47:33-8.
11. Streubel A, Siepmann J, Dashevsky A, Bodmeier R. pH-independent release of a weakly basic drug from water-insoluble and -soluble matrix tablets. *J. Control. Release* 2000;67:101-10.
12. Rubinstein MH, Garr JSM. Compaction properties of a cellulose-lactose direct-compression excipient. *Pharm. Technol. Int.* 1991;3:24-7.
13. Fernández-Arévalo M, Vela MT, Rabasco AM. Rheological study of lactose coated with acrylic resins. *Drug Dev. Ind. Pharm.* 1990;16:295-313.
14. Ho HO, Wang, HY, Sheu MT. The evaluation of granulated excipients as matrix materials for controlled delivery of captopril. *J. Control. Release*, 1997;49:243-51.
15. Carr RL. Evaluating flow properties of solids. *Chem. Eng.* 1965;72:163-8.
16. Tsai T, San YP, Ho HO, Kao CC, Sheu MT. Film-forming polymer granulated excipients as matrix materials for the production of pellets by extrusion/spheronization. *Chin. Pharm. J.* 1997;49:249-57.
17. Tsai T, San YP, Ho HO, Wu JS, Sheu MT. Film-forming polymer-granulated excipients as the matrix materials for controlled release dosage forms. *J. Control. Release* 1998;51:289-99.
18. Makagawa H. "Application of solid dispersions to prepare some oral dosage forms" in *Advances in pharmaceuticals and pharmaceutical technology*. S.Y. Lin Eds.; Jeou Chou Book. Taipei, Taiwan, 1989; pp 203-11.