

SciFinder - Explore References - Windows Internet Explorer

https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf

SciFinder®

Welcome ChiHua Lu | Sign Out

Add KMP Alert Research Topic "amphetamine" > references (45890)

Explore References

Research Topic: Search

Examples:
The effect of antibiotic residues on dairy products
Photocyanation of aromatic compounds

Author Name
Company Name
Document Identifier
Journal
Patent
Tags

Publication Year(s)
Examples: 1995, 1995-1999, 1995-, -1995

Document Type(s)

☐ Biography	☐ Dissertation	☐ Patent
☐ Book	☐ Editorial	☐ Preprint
☐ Clinical Trial	☐ Historical	☐ Report
☐ Commentary	☐ Journal	☐ Review
☐ Conference	☐ Letter	

Language(s)

☐ Chinese	☐ German	☐ Polish
----------------------------------	---------------------------------	---------------------------------

Saved Answer Sets

Autosaved Reference Set

View All

Import

Keep Me Posted Results

No profiles exist

完成

SciFinder - Reference Answer Set - Windows Internet Explorer

https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf

SciFinder - Reference Answer Set

References

167 References 0 Selected

Select All Deselect All Sort by: Accession Number

Answers per Page [20] 1 2 3 4 5 6 ... 9

Display: - +

1. The effect of weight loss and treatment with metformin on serum vaspin levels in women with polycystic ovary syndrome

By Koio, Ekaterini Tzomalos, Konstantinos; Dinas, Konstantinos; Katsikis, Ilias; Kalaitzakis, Emmanuil; Delkos, Dimitrios; Kandaraki, Eleni A.; Panidis, Dimitrios

From Endocrine Journal (Kyoto, Japan) (2011), 58(4), 237-246. Language: English, Database: CAPLUS

Many patients with polycystic ovary syndrome (PCOS) have insulin resistance, obesity (mostly visceral) and glucose intolerance, conditions assocd. with abnormalities in the prodn. of vaspin, a novel adipokine that appears to preserve insulin sensitivity and glucose tolerance. The aim of the study was to assess serum vaspin levels in PCOS and the effects on vaspin levels of metformin or of wt. loss. We studied 79 patients with PCOS and 50 healthy female volunteers. Normal wt. patients with PCOS (n = 25) were treated with metformin 850 mg bid for 6 mo. Overweight/obese patients with PCOS (n ...

+Substances Reactions ~1 Citing Full Text Link 0 Comments 0 Tags

2. Microscopic trends in methanol/water and acetonitrile/water systems

By Alzweiri, Muhammed; Parkinson, John A.; Watson, David W.; Steer, Steve

From Jordan Journal of Pharmaceutical Sciences (2011), 4(1), 20-28. Language: English, Database: CAPLUS

The most practical measure of therapeutic equivalence between two com. available and generic formulation of a certain drug is to det. their in vivo bioavailability. However, for the oral dosage form that is not intended to be absorbed (e.g. orlistat), in vivo bioavailability studies are irrelevant to the achievement of the product's intended purposes. However, specific requirements for these drug products may be set in a way that they should meet acceptable in vitro stds. For this purpose, a comparative enzymic inhibition assay of the target enzyme, pancreatic lipase, was developed to demon...

+Substances Reactions ~0 Citings Full Text Link 0 Comments 0 Tags

3. Evaluation of carcinogenicity studies of medicinal products for human use authorised via the European centralised procedure (1995-2009)

By Friedrich, Anita; Olejniczak, Klaus

From Regulatory Toxicology and Pharmacology (2011), 60(2), 225-248. Language: English, Database: CAPLUS

Carcinogenicity data of medicinal products for human use that have been authorised via the European centralised procedure (CP) between 1995 and 2009 were evaluated. Carcinogenicity data, either from long-term rodent carcinogenicity studies, transgenic mouse studies or repeat-dose toxicity studies were available for 144 active

Analysis Refine

Analyze by:

Author Name

Click bar to view only those references within the current answer set

Anonymous	6
Scheen A J	5
Bray George A	4
Hauptman Jonathan	4
Richardson Robyn D	4
Romo Daniel	4
Smith Jeffrey W	4
Boss Olivier	3
Hofbauer Karl G	3
Rissanen Aila	3

Show More

Categorize

完成

The effect of weight loss and treatment with metformin on serum vaspin levels in women with polycystic ovary syndrome



By: Koïou, Ekaterini; Tziomalos, Konstantinos; Dinas, Konstantinos; Katsikis, Ilias; Kalaitzakis, Emmanuil; Delkos, Dimitrios; Kandaraki, Eleni A.; Panidis, Dimitrios

Many patients with polycystic ovary syndrome (PCOS) have insulin resistance, obesity (mostly visceral) and glucose intolerance, conditions assocd. with abnormalities in the prodn. of vaspin, a novel adipokine that appears to preserve insulin sensitivity and glucose tolerance. The aim of the study was to assess serum vaspin levels in PCOS and the effects on vaspin levels of metformin or of wt. loss. We studied 79 patients with PCOS and 50 healthy female volunteers. Normal wt. patients with PCOS (n = 25) were treated with metformin 850 mg bid for 6 mo. Overweight/obese patients with PCOS (n = 54) were prescribed a normal-protein, energy-restricted diet for 6 mo; half of them were also given orlistat 120 mg tid and the rest were given sibutramine 10 mg qd. At baseline and after 6 mo, serum vaspin levels and anthropometric, metabolic and hormonal features of PCOS were detd. Overall, patients with PCOS had higher vaspin levels than controls (p = 0.021). Normal wt. patients with PCOS had higher vaspin levels than normal wt. controls (p = 0.043). Vaspin levels were non-significantly higher in overweight/obese patients with PCOS than in overweight/obese controls. In normal wt. patients with PCOS, metformin reduced vaspin levels non-significantly. In overweight/obese patients with PCOS, diet plus orlistat or sibutramine did not affect vaspin levels. Vaspin levels were independently correlated with body mass index in women with PCOS (p = 0.001) and with waist circumference in controls (p = 0.015). In conclusion, serum vaspin levels are elevated in PCOS but neither a small wt. loss nor metformin affect vaspin levels significantly.

心得：

在 Scifinder 上，有數以萬計的資料可供我們找尋，也可以把未知的結構化在上面，尋找看看是否有前輩或其他國家學者已經有研究過這個結構，對於研究生及做研究的學者來說，這是個必備的軟體，也是必備的能力。

對於我們新手來說，一個很大的障礙也是好處的地方是，他全是英文!!經過這次作業下來，大概又加強了英文能力吧。不過學校使用一次只能兩個人上線有點太少了，這麼好用的軟體應該會供不應求吧。



Thomson Innovation - Windows Internet Explorer

http://www.thomsoninnovation.com/hip-innovation/recordView.do?pager.offset=0&recordCount=1&parentId=null&totalRecords=1815&fromRecordView=false&dataSource=T3&category=PAT&selectedRecord=

Patent Record View - US8080526B2 [Close Record View](#)

Record View: Patent US8080526B2 [Help](#)

[Add to Work File](#) | [Mark Record](#) | [Watch Record](#) | [Download](#) | [Print](#)

Quick View [Go to Full View](#)

DWPI Title ?

Original Title ?
Methods and compositions for the treatment of gastrointestinal disorders


Abstract ?
The present invention features compositions and related methods for treating IBS and other gastrointestinal disorders and conditions (e.g., gastrointestinal motility disorders, functional gastrointestinal disorders, gastroesophageal reflux disease (GERD), Crohn's disease, ulcerative colitis, Inflammatory bowel disease, functional heartburn, dyspepsia (including functional dyspepsia or nonulcer dyspepsia), gastroparesis, chronic intestinal pseudo-obstruction (or colonic pseudo-obstruction), and disorders and conditions associated with constipation, e.g., constipation associated with use of opiate pain killers, post-surgical constipation, and constipation associated with neuropathic disorders as well as other conditions and disorders using peptides and other agents that activate the guanylate cyclase C (GC-C) receptor.

First Claim ?
1. A peptide consisting of the amino acid sequence Cys6 Cys7 Glu8 Tyr9 Cys10 Cys11 Asn12 Pro13 Ala14 Cys15 Thr16 Gly17 Cys18 Tyr19 (SEQ ID NO: 31), wherein disulfide bonds are present between Cys6-Cys11, Cys7-Cys15, and Cys10-Cys18.

Assignee / Applicant ?

Record 1 of 1815

Images [Image 1/19](#) [Zoom \(+\)](#)



[Mark and Next](#)

完成

開始 日投第3年... 新增Microso... Login | Thom... Thomson Inn... Thomson Inn... CH 100% 上午 11:51

Abstract

?

The present invention features compositions and related methods for treating IBS and other gastrointestinal disorders and conditions (e.g., gastrointestinal motility disorders, functional gastrointestinal disorders, gastroesophageal reflux disease (GERD), Crohn's disease, ulcerative colitis, Inflammatory bowel disease, functional heartburn, dyspepsia (including functional dyspepsia or nonulcer dyspepsia), gastroparesis, chronic intestinal pseudo-obstruction (or colonic pseudo-obstruction), and disorders and conditions associated with constipation, e.g., constipation associated with use of opiate pain killers, post-surgical constipation, and constipation associated with neuropathic disorders as well as other conditions and disorders using peptides and other agents that activate the guanylate cyclase C (GC-C) receptor.

心得：

專利檢索平台的優點在，他時時刻刻都在更新，而且是彷彿無國界般，大部份每個國家的專利都有收錄並翻譯成英文，這樣可以省去很多時間。

專利不只是為人類創造福利，也是創造財富的一項資產，如果以後有志向開公司成立企業的人，這套軟體是很有用處的，特別是在國際上，有許許多多的專利限定，常常有藥的結構是因為要避專利所以添加另外一些結構，專利權在我們身邊無所不在。